

ELLIPTIC BOUNDARY-VALUE PROBLEMS IN THE SENSE OF LAWРУK ON SOBOLEV AND HÖRMANDER SPACES

ЕЛІПТИЧНІ КРАЙОВІ ЗАДАЧІ ЗА ЛАВРУКОМ У ПРОСТОРАХ СОБОЛЄВА І ХЕРМАНДЕРА

We study elliptic boundary-value problems with additional unknown functions in boundary conditions. These problems were introduced by Lawruk. We prove that the operator corresponding to a problem of this kind is bounded and Fredholm in appropriate couples of the inner product isotropic Hörmander spaces $H^{s,\varphi}$, which form the refined Sobolev scale. The order of differentiation for these spaces is given by a real number s and a positive function φ slowly varying at infinity in Karamata's sense. We consider this problem for an arbitrary elliptic equation $Au = f$ in a bounded Euclidean domain Ω under the condition that $u \in H^{s,\varphi}(\Omega)$, $s < \text{ord } A$, and $f \in L_2(\Omega)$. We prove theorems on the *a priori* estimate and regularity of the generalized solutions to this problem.

Досліджено еліптичну крайову задачу з додатковими невідомими функціями у крайових умовах. Ці задачі введені Лавруком. Доведено, що оператор, відповідний такій задачі, є обмеженим і нетеровим у відповідних парах гільбертових ізотропних просторів Хермандера $H^{s,\varphi}$, які утворюють уточнену соболевську шкалу. Показник диференційовності для цих просторів задано дійсним числом s і додатною функцією φ , яка повільно змінюється на нескінченності за Караматою. Ця задача розглядається для довільного еліптичного рівняння $Au = f$ в евклідовій області Ω за умов, що $u \in H^{s,\varphi}(\Omega)$, $s < \text{ord } A$ і $f \in L_2(\Omega)$. Доведено теореми про апіорну оцінку і регулярність узагальнених розв'язків цієї задачі.

1. Introduction. In this paper, we investigate elliptic boundary-value problems with additional unknown functions in boundary conditions. These functions are given on the smooth boundary of a Euclidean domain Ω , where the problem is posed. Such problems were introduced by B. Lawruk [1–3]. They appear naturally if we pass from a general (nonregular) elliptic boundary-value problem to its formally adjoint problem. Moreover, the class of these problems are closed with respect to this passage. Their various examples occur in hydrodynamics and the theory of elasticity [4–6].

This class has been investigated in some function spaces by V. A. Kozlov, V. G. Maz'ya, and J. Rossmann [7] (Chapt. 3), mainly, for scalar elliptic equations and by I. Ya. Roitberg [8, 9] for mixed-order elliptic systems. The results of I. Ya. Roitberg is expounded in Ya. A. Roitberg's monograph [10] (Chapt. 2). The main result obtained are the theorem on the Fredholm property of these problems on two-sided scales of normed spaces, solutions of the elliptic equations being considered in the spaces introduced by Ya. A. Roitberg [11, 12] (see also his monograph [13] (Sect. 2)). These spaces coincide with the Sobolev spaces provided that their differentiation order is large enough. But, for other orders, the Roitberg spaces contain elements that are not distributions on Ω .

The purpose of this paper is to prove a version of this theorem for Sobolev and more general Hörmander spaces [14] (Sect. 2.2) of distributions on Ω . We use the Hörmander spaces

$$H^{s,\varphi}(\mathbb{R}^n) := \{w \in \mathcal{S}'(\mathbb{R}^n) : \langle \xi \rangle^s \varphi(\langle \xi \rangle) \widehat{w}(\xi) \in L_2(\mathbb{R}^n, d\xi)\}$$

and their analogs for the domain Ω and its boundary. These spaces are parametrized with the number $s \in \mathbb{R}$ and the function $\varphi : [1, \infty) \rightarrow (0, \infty)$ that varies slowly at infinity in the sense of J. Karamata.

Here, \widehat{w} is the Fourier transform of the tempered distribution w , whereas $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$. If $\varphi \equiv 1$, then $H^{s,\varphi}(\mathbb{R}^n)$ becomes the Sobolev space $H^s(\mathbb{R}^n)$.

These Hörmander spaces form the refined Sobolev scale, which was selected and investigated by V. A. Mikhailets and A. A. Murach [15–17]. This scale has an important interpolation property; namely, every space $H^{s,\varphi}(\mathbb{R}^n)$ is a result of the interpolation with an appropriate function parameter of the inner product Sobolev spaces $H^{s-\varepsilon}(\mathbb{R}^n)$ and $H^{s+\delta}(\mathbb{R}^n)$ with $\varepsilon, \delta > 0$.

V. A. Mikhailets and A. A. Murach [15, 16, 18–23] have elaborated the theory of solvability of general elliptic boundary-value problems on the refined Sobolev scale and its modification by Ya. A. Roitberg. The above-mentioned interpolation is a key method in their theory. In this connection, we also note papers [24–26]. However, this theory does not involve the important class of elliptic boundary-value problems in the sense of B. Lawruk.

In this paper, we consider these problems for an arbitrary scalar elliptic equation $Au = f$ with $u \in H^{s,\varphi}(\Omega)$ for $s < \text{ord } A$ and $f \in L_2(\Omega)$. This approach originates from the papers by J.-L. Lions and E. Magenes [27, 28] (see also [29]). The case of $s \geq \text{ord } A$ was investigated in [30].

This paper consists of seven sections. In Section 2, we formulate an elliptic boundary-value problem in the sense of B. Lawruk and discuss the corresponding Green formula and formally adjoint problem. In Section 3, we give the definitions of Hörmander function spaces that form the refined Sobolev scale. Section 4 contains the main results of the paper. They are the theorem on the Fredholm property of the problem under investigation, the a priori estimate for its generalized solutions, and the theorem about their regularity on the refined Sobolev scale. In Sections 5 and 6, we discuss auxiliary results that we need to prove these theorems. The proofs of the main results are given in Section 7.

2. Statement of the problem. Let Ω be a bounded domain in \mathbb{R}^n , where $n \geq 2$. We suppose that its boundary $\Gamma := \partial\Omega$ is a closed C^∞ -manifold of dimension $n - 1$, the C^∞ -structure on Γ being induced by \mathbb{R}^n . Let $\nu(x)$ denote the unit vector of the inward normal to Γ at a point $x \in \Gamma$.

Choose arbitrarily integers $q \geq 1$, $\varkappa \geq 1$, $m_1, \dots, m_{q+\varkappa} \in [0, 2q - 1]$ and r_1, \dots, r_\varkappa . In the paper, we consider the linear boundary-value problem on the domain Ω with \varkappa additional unknown functions on the boundary Γ :

$$Au = f \quad \text{on } \Omega, \quad (1)$$

$$B_j u + \sum_{k=1}^{\varkappa} C_{j,k} v_k = g_j \quad \text{on } \Gamma, \quad j = 1, \dots, q + \varkappa. \quad (2)$$

Here,

$$A := A(x, D) := \sum_{|\mu| \leq 2q} a_\mu(x) D^\mu$$

is a linear differential operator on $\overline{\Omega} := \Omega \cup \Gamma$ of the even order $2q$. Besides, each

$$B_j(x, D) := \sum_{|\mu| \leq m_j} b_{j,\mu}(x) D^\mu$$

is a linear boundary differential operator on Γ of order m_j , and every $C_{j,k} := C_{j,k}(x, D_\tau)$ is a linear tangent differential operator on Γ with $\text{ord } C_{j,k} \leq m_j + r_k$. All the coefficients of these

differential operators are supposed to be infinitely smooth complex-valued functions given on $\bar{\Omega}$ and Γ respectively.

Here and below, we use the following standard notation: $\mu := (\mu_1, \dots, \mu_n)$ is a multiindex with $|\mu| := \mu_1 + \dots + \mu_n$, and $D^\mu := D_1^{\mu_1} \dots D_n^{\mu_n}$ with $D_\ell := i\partial/\partial x_\ell$, where i is imaginary unit and $x = (x_1, \dots, x_n) \in \mathbb{R}^n$. Besides, $D_\nu := i\partial/\partial \nu(x)$ and $\xi^\mu := \xi_1^{\mu_1} \dots \xi_n^{\mu_n}$ for $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{C}^n$.

The function u on Ω and all the functions v_1, \dots, v_\varkappa on Γ are unknown in the problem (1), (2). In the paper, all functions and distributions are supposed to be complex-valued.

We assume that the boundary-value problem (1), (2) is elliptic on Ω in the sense of B. Lawruk [1]. Recall the corresponding definition (see also [7] (Subsect. 3.1.3)).

Let $A^{(0)}(x, \xi)$ and $B_j^{(0)}(x, \xi)$ denote the principal symbols of the differential operators $A(x, D)$ and $B_j(x, D)$ respectively. We recall that

$$A^{(0)}(x, \xi) := \sum_{|\mu|=2q} a_\mu(x) \xi^\mu, \quad \text{with } x \in \bar{\Omega} \quad \text{and } \xi \in \mathbb{C}^n,$$

is a homogeneous polynomial of order $2q$ in ξ and that

$$B_j^{(0)}(x, \xi) := \sum_{|\mu|=m_j} b_{j,\mu}(x) \xi^\mu, \quad \text{with } x \in \Gamma \quad \text{and } \xi \in \mathbb{C}^n,$$

is a homogeneous polynomial of order m_j in ξ . Besides, let $C_{j,k}^{(0)}(x, \tau)$ denote the principal symbol of $C_{j,k}(x, D_\tau)$ if $\text{ord } C_{j,k} = m_j + r_k$. For every point $x \in \Gamma$, the expression $C_{j,k}^{(0)}(x, \tau)$ is a homogeneous polynomial of order $m_j + r_k$ in τ , with τ being a tangent vector to the boundary Γ at the point x . (If $\text{ord } C_{j,k} < m_j + r_k$, then we put $C_{j,k}^{(0)}(x, \tau) := 0$.)

The boundary-value problem (1), (2) is said to be elliptic on Ω if the following three conditions are fulfilled:

(i) The differential operator $A(x, D)$ is elliptic at each point $x \in \bar{\Omega}$; i.e., $A^{(0)}(x, \xi) \neq 0$ for an arbitrary vector $\xi \in \mathbb{R}^n \setminus \{0\}$.

(ii) The differential operator $A(x, D)$ is properly elliptic at each point $x \in \Gamma$; i.e., for an arbitrary vector $\tau \neq 0$ tangent to Γ at x , the polynomial $A^{(0)}(x, \tau + \zeta \nu(x))$ in $\zeta \in \mathbb{C}$ has q roots with positive imaginary part and q roots with negative imaginary part (these roots are calculated with regard for their multiplicity).

(iii) The system of boundary-value conditions (2) covers equation (1) at each point $x \in \Gamma$. This means that, for an arbitrary vector $\tau \neq 0$ from condition (ii), the boundary-value problem

$$A^{(0)}(x, \tau + D_t \nu(x)) \theta(t) = 0 \quad \text{for } t > 0,$$

$$B_j^{(0)}(x, \tau + D_t \nu(x)) \theta(t) \Big|_{t=0} + \sum_{k=1}^{\varkappa} C_{j,k}^{(0)}(x, \tau) \lambda_k = 0, \quad j = 1, \dots, q + \varkappa,$$

has only zero solution. This problem is considered relative to the unknown function $\theta \in C^\infty([0, \infty))$ with $\theta(t) \rightarrow 0$ as $t \rightarrow \infty$ and the unknown complex-valued numbers $\lambda_1, \dots, \lambda_\varkappa$. Here, $A^{(0)}(x, \tau + D_t \nu(x))$ and $B_j^{(0)}(x, \tau + D_t \nu(x))$ are differential operators with respect to $D_t := i\partial/\partial t$. We obtain them if we put $\zeta := D_t$ in the polynomials $A^{(0)}(x, \tau + \zeta \nu(x))$ and $B_j^{(0)}(x, \tau + \zeta \nu(x))$ in ζ .

It is worthwhile to note that condition (ii) follows from condition (i) in the $n \geq 3$ case.

Examples of elliptic boundary-value problems in the sense of B. Lawruk are discussed in [7] (Subsect. 3.1.5). Among them, we mention the following simple example:

$$\Delta u = f \quad \text{on } \Omega, \quad u + v = g_1 \quad \text{and} \quad D_\nu u + D_\tau v = g_2 \quad \text{on } \Gamma.$$

Here, $n = 2$, $\varkappa = 1$, Δ is the Laplace operator, and $D_\tau := i\partial/\partial\tau$, with $\partial/\partial\tau$ being the derivative along the curve Γ . It is easy to check that this boundary-value problem is elliptic on Ω .

To the problem (1), (2) corresponds the linear mapping

$$\Lambda: (u, v_1, \dots, v_\varkappa) \mapsto \left(Au, B_1 u + \sum_{k=1}^{\varkappa} C_{1,k} v_k, \dots, B_{q+\varkappa} u + \sum_{k=1}^{\varkappa} C_{q+\varkappa,k} v_k \right), \quad (3)$$

$$\text{with } u \in C^\infty(\bar{\Omega}) \quad \text{and} \quad v_1, \dots, v_\varkappa \in C^\infty(\Gamma).$$

We will investigate properties of the extension (by continuity) of this mapping in suitable couples of Hörmander function spaces and, in particular, Sobolev spaces.

In order to describe the range of this extension, we need the following Green formula (see [7] (Theorem 3.1.1)):

$$\begin{aligned} & (Au, w)_\Omega + \sum_{j=1}^{q+\varkappa} \left(B_j u + \sum_{k=1}^{\varkappa} C_{j,k} v_k, h_j \right)_\Gamma = \\ & = (u, A^+ w)_\Omega + \sum_{j=1}^{2q} \left(D_\nu^{j-1} u, K_j w + \sum_{k=1}^{q+\varkappa} Q_{k,j}^+ h_k \right)_\Gamma + \sum_{j=1}^{\varkappa} \left(v_j, \sum_{k=1}^{q+\varkappa} C_{k,j}^+ h_k \right)_\Gamma, \end{aligned}$$

where $u, w \in C^\infty(\bar{\Omega})$, $v = (v_1, \dots, v_\varkappa) \in (C^\infty(\Gamma))^\varkappa$, and $h = (h_1, \dots, h_{q+\varkappa}) \in (C^\infty(\Gamma))^{q+\varkappa}$ are arbitrary, whereas $(\cdot, \cdot)_\Omega$ and $(\cdot, \cdot)_\Gamma$ are the inner products in the Hilbert spaces $L_2(\Omega)$ and $L_2(\Gamma)$ of functions square integrable over Ω and Γ respectively.

Here, A^+ stands for the differential operator that is formally adjoint to A with respect to $(\cdot, \cdot)_\Omega$. Besides, $C_{k,j}^+$ and $Q_{k,j}^+$ denote the tangent differential operators that are formally adjoint to $C_{k,j}$ and $Q_{k,j}$ respectively relative to $(\cdot, \cdot)_\Gamma$, the tangent differential operators $Q_{k,j}$ appearing in the representation of the boundary differential operators B_j in the form

$$B_j(x, D) = \sum_{k=1}^{2q} Q_{j,k}(x, D_\tau) D_\nu^{k-1}.$$

(If $k > m_j$, then $Q_{j,k} = 0$.) Finally, $K_j := K_j(x, D)$ is a certain linear boundary differential operator on Γ , with $\text{ord } K_j \leq 2q - j$.

Taking into account the Green formula, we consider the following boundary-value problem on Ω with $q + \varkappa$ additional unknown function on Γ :

$$A^+ w = \omega \quad \text{in } \Omega, \quad (4)$$

$$K_j w + \sum_{k=1}^{q+\varkappa} Q_{k,j}^+ h_k = \chi_j \quad \text{on } \Gamma, \quad j = 1, \dots, 2q, \quad (5)$$

$$\sum_{k=1}^{q+\varkappa} C_{k,j}^+ h_k = \chi_{2q+j} \quad \text{on } \Gamma, \quad j = 1, \dots, \varkappa. \quad (6)$$

This problem is formally adjoint to the problem (1), (2) with respect to the Green formula. It is worthwhile to note that the problem (1), (2) is elliptic if and only if the problem (4)–(6) is elliptic as well (see [7] (Theorem 3.1.2)).

3. Hörmander spaces and refined Sobolev scales. Here, we give the definitions of the Hörmander inner product spaces that form the refined Sobolev scales over \mathbb{R}^n , Ω , and Γ . We also discuss some of their properties [23].

We should begin with the definition of the Hörmander spaces $H^{s,\varphi}(\mathbb{R}^n)$, with $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. Here and below, \mathcal{M} stands for the set of all Borel measurable functions $\varphi: [1, \infty) \rightarrow (0, \infty)$ such that both the functions φ and $1/\varphi$ are bounded on each compact interval $[1, b]$, with $1 < b < \infty$, and that the function φ is slowly varying at infinity in the sense of J. Karamata [31]. The latter property means that $\varphi(\lambda t)/\varphi(t) \rightarrow 1$ as $t \rightarrow \infty$ for every $\lambda > 0$.

Note that the slowly varying functions are well investigated and has various applications (see monographs [32, 33]). We obtain an important example of these functions by setting

$$\varphi(t) := (\log t)^{r_1} (\log \log t)^{r_2} \dots \underbrace{(\log \dots \log t)^{r_k}}_{k \text{ times}} \quad \text{of } t \gg 1.$$

Here, $k \in \mathbb{Z}$, with $k \geq 1$, and $r_1, \dots, r_k \in \mathbb{R}$ are arbitrary parameters.

Let $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. By definition, the complex linear space $H^{s,\varphi}(\mathbb{R}^n)$, with $n \geq 1$, consists of all distributions $w \in \mathcal{S}'(\mathbb{R}^n)$ that the Fourier transform \widehat{w} of w is locally Lebesgue integrable over \mathbb{R}^n and satisfies the condition

$$\int_{\mathbb{R}^n} \langle \xi \rangle^{2s} \varphi^2(\langle \xi \rangle) |\widehat{w}(\xi)|^2 d\xi < \infty.$$

Here, as usual, $\mathcal{S}'(\mathbb{R}^n)$ is the linear topological space of all tempered distributions on \mathbb{R}^n , and $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$ is the smoothed modulus of $\xi \in \mathbb{R}^n$. The space $H^{s,\varphi}(\mathbb{R}^n)$ is endowed with the inner product

$$(w_1, w_2)_{H^{s,\varphi}(\mathbb{R}^n)} := \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} \varphi^2(\langle \xi \rangle) \widehat{w}_1(\xi) \overline{\widehat{w}_2(\xi)} d\xi.$$

It naturally induces the norm

$$\|w\|_{H^{s,\varphi}(\mathbb{R}^n)} := (w, w)_{H^{s,\varphi}(\mathbb{R}^n)}^{1/2}.$$

The space $H^{s,\varphi}(\mathbb{R}^n)$ just defined is a special isotropic Hilbert case of the spaces $\mathcal{B}_{p,\mu}$ introduced and investigated by L. Hörmander in [14] (Sect. 2.2) (see also his monograph [34] (Sect. 10.1)). Namely, $H^{s,\varphi}(\mathbb{R}^n) = \mathcal{B}_{p,\mu}$ if $p = 2$ and $\mu(\xi) \equiv \langle \xi \rangle^s \varphi(\langle \xi \rangle)$. Note that the inner product spaces $\mathcal{B}_{2,\mu}$ were also investigated by L. R. Volevich and B. P. Paneah in [35] (§ 2).

In the $\varphi \equiv 1$ case, the space $H^{s,\varphi}(\mathbb{R}^n)$ becomes the inner product Sobolev space $H^s(\mathbb{R}^n)$ of order $s \in \mathbb{R}$. Generally, we have the continuous and dense embeddings

$$H^{s+\varepsilon}(\mathbb{R}^n) \hookrightarrow H^{s,\varphi}(\mathbb{R}^n) \hookrightarrow H^{s-\varepsilon}(\mathbb{R}^n) \quad \text{for every } \varepsilon > 0. \quad (7)$$

They show that, the numerical parameter s characterizes the main regularity of the distributions $w \in H^{s,\varphi}(\mathbb{R}^n)$, whereas the function parameter φ defines a supplementary regularity, which is

subordinate to s . Specifically, if $\varphi(t) \rightarrow \infty$ [or $\varphi(t) \rightarrow 0$] as $t \rightarrow \infty$, then the parameter φ defines the supplementary positive [or negative] regularity. Thus, we can say that φ refines the main regularity s in the class of Hilbert separable spaces

$$\{H^{s,\varphi}(\mathbb{R}^n) : s \in \mathbb{R}, \varphi \in \mathcal{M}\}.$$

This class is selected by V. A. Mikhailets and A. A. Murach in [15] and is called the refined Sobolev scale over \mathbb{R}^n (see also monograph [23] (Subsect. 1.3.3)).

In the paper, we need the refined Sobolev scales over the domain Ω and its boundary Γ . These scales are constructed in the standard way with the help of the Hörmander spaces $H^{s,\varphi}(\mathbb{R}^n)$. Let us pass to the corresponding definitions [23] (Sect. 2.1 and 3.1).

The complex linear space $H^{s,\varphi}(\Omega)$ and the norm in it are defined as follows:

$$H^{s,\varphi}(\Omega) := \{w \upharpoonright \Omega : w \in H^{s,\varphi}(\mathbb{R}^n)\},$$

$$\|u\|_{H^{s,\varphi}(\Omega)} := \inf \{\|w\|_{H^{s,\varphi}(\mathbb{R}^n)} : w \in H^{s,\varphi}(\mathbb{R}^n), u = w \upharpoonright \Omega\},$$

with $u \in H^{s,\varphi}(\Omega)$. Here, as usual, $w \upharpoonright \Omega$ stands for the restriction of the distribution w to the domain Ω . The space $H^{s,\varphi}(\Omega)$ is Hilbert and separable with respect to this norm. The space $H^{s,\varphi}(\Omega)$ is continuously embedded in the linear topological space $\mathcal{S}'(\Omega) := \{w \upharpoonright \Omega : w \in \mathcal{S}'(\mathbb{R}^n)\}$. The set $C^\infty(\overline{\Omega})$ is dense in $H^{s,\varphi}(\Omega)$.

In short, the space $H^{s,\varphi}(\Gamma)$ consists of all distributions on Γ that belong, in local coordinates, to $H^{s,\varphi}(\mathbb{R}^{n-1})$. Let us dwell on the definition of $H^{s,\varphi}(\Gamma)$. From C^∞ -structure on Γ , we arbitrarily choose a finite collection of local charts $\alpha_j : \mathbb{R}^{n-1} \leftrightarrow \Gamma_j$, with $j = 1, \dots, \lambda$, that the open sets $\Gamma_1, \dots, \Gamma_\lambda$ form a covering of the manifold Γ . Besides, we arbitrarily choose functions $\chi_j \in C^\infty(\Gamma)$, with $j = 1, \dots, \lambda$, that form a partition of unity on Γ which satisfies the condition $\text{supp } \chi_j \subset \Gamma_j$. Then the complex linear space $H^{s,\varphi}(\Gamma)$ and the norm in it are defined as follows:

$$H^{s,\varphi}(\Gamma) := \{h \in \mathcal{D}'(\Gamma) : (\chi_j h) \circ \alpha_j \in H^{s,\varphi}(\mathbb{R}^{n-1}) \text{ for every } j \in \{1, \dots, \lambda\}\},$$

$$\|h\|_{H^{s,\varphi}(\Gamma)} := \left(\sum_{j=1}^{\lambda} \|(\chi_j h) \circ \alpha_j\|_{H^{s,\varphi}(\mathbb{R}^{n-1})}^2 \right)^{1/2}.$$

Here, as usual, $\mathcal{D}'(\Gamma)$ is the linear topological space of all distributions on Γ , whereas $(\chi_j h) \circ \alpha_j$ is the representation of the distribution $\chi_j h$ in the local chart α_j . The space $H^{s,\varphi}(\Gamma)$ is Hilbert and separable with respect to this norm. This space does not depend (up to equivalence of norms) on the indicated choice of the local charts and partition of unity [23] (Theorem 2.3). The space $H^{s,\varphi}(\Gamma)$ is continuously embedded in $\mathcal{D}'(\Gamma)$. The set $C^\infty(\Gamma)$ is dense in $H^{s,\varphi}(\Gamma)$.

Thus, we have the refined Sobolev scales

$$\{H^{s,\varphi}(\Omega) : s \in \mathbb{R}, \varphi \in \mathcal{M}\} \quad \text{and} \quad \{H^{s,\varphi}(\Gamma) : s \in \mathbb{R}, \varphi \in \mathcal{M}\} \quad (8)$$

over Ω and Γ respectively. They contain the inner product Sobolev spaces $H^s(\Omega) := H^{s,1}(\Omega)$ and $H^s(\Gamma) := H^{s,1}(\Gamma)$. (Here, of course, 1 denotes the function equaled identically to 1). It follows from (7) that

$$H^{s+\varepsilon}(\Omega) \hookrightarrow H^{s,\varphi}(\Omega) \hookrightarrow H^{s-\varepsilon}(\Omega) \quad \text{for every } \varepsilon > 0, \quad (9)$$

$$H^{s+\varepsilon}(\Gamma) \hookrightarrow H^{s,\varphi}(\Gamma) \hookrightarrow H^{s-\varepsilon}(\Gamma) \quad \text{for every } \varepsilon > 0. \quad (10)$$

These embeddings are compact and dense.

It is worthwhile to note the following connection between the scales (8) (see [23] (Subsect. 3.2.1)). Let $s > 1/2$ and $\varphi \in \mathcal{M}$, then the trace mapping $u \mapsto u|_{\Gamma}$, with $u \in C^\infty(\Gamma)$, extends uniquely (by continuity) to a bounded surjective operator $R_\Gamma: H^{s,\varphi}(\Omega) \rightarrow H^{s-1/2,\varphi}(\Gamma)$. Thus, for each distribution $u \in H^{s,\varphi}(\Omega)$, its trace $R_\Gamma u$ on Γ is well defined. Moreover, we have the equivalence of norms

$$\|h\|_{H^{s-1/2,\varphi}(\Gamma)} \asymp \inf \{ \|u\|_{H^{s,\varphi}(\Omega)} : u \in H^{s,\varphi}(\Omega), h = R_\Gamma u \}$$

on the class of all $h \in H^{s-1/2,\varphi}(\Gamma)$. But it is impossible to define this trace reasonably in the $s < 1/2$ case.

4. Main results. To formulate them, we need to introduce the linear spaces N and N^+ of infinitely smooth solutions to the problem (1), (2) and its formally adjoint problem (4)–(6). Namely, let N consist of all solutions

$$(u, v_1, \dots, v_\varkappa) \in C^\infty(\bar{\Omega}) \times (C^\infty(\Gamma))^\varkappa$$

to the problem (1), (2) in the case where $f = 0$ on Ω and all $g_j = 0$ on Γ . Besides, let N^+ consist of all solutions

$$(w, h_1, \dots, h_{q+\varkappa}) \in C^\infty(\bar{\Omega}) \times (C^\infty(\Gamma))^{q+\varkappa}$$

to the problem (4)–(6) in the case where $\omega = 0$ on Ω and all $\chi_j = 0$ and all $\chi_{2q+j} = 0$ on Γ . Since these problems are elliptic on Ω , the spaces N and N^+ are finite-dimensional [7] (Lemma 3.4.2).

We are motivated by the following result.

Proposition 1. *For arbitrary $s > 2q - 1/2$ and $\varphi \in \mathcal{M}$, the mapping (3) extends uniquely (by continuity) to a bounded operator*

$$\Lambda: H^{s,\varphi}(\Omega) \oplus \bigoplus_{k=1}^{\varkappa} H^{s+r_k-1/2,\varphi}(\Gamma) \rightarrow H^{s-2q,\varphi}(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{s-m_j-1/2,\varphi}(\Gamma). \quad (11)$$

This operator is Fredholm. Its kernel coincides with N , and its range consists of all vectors

$$(f, g_1, \dots, g_{q+\varkappa}) \in H^{s-2q,\varphi}(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{s-m_j-1/2,\varphi}(\Gamma)$$

such that

$$(f, w)_\Omega + \sum_{j=1}^{q+\varkappa} (g_j, h_j)_\Gamma = 0 \quad \text{for all } (w, h_1, \dots, h_{q+\varkappa}) \in N^+. \quad (12)$$

The index of the operator (11) is equal to $\dim N - \dim N^+$ and does not depend on s and φ .

Here and below, $(\cdot, \cdot)_\Omega$ and $(\cdot, \cdot)_\Gamma$ denote the extension by continuity of the inner products in the Hilbert spaces $L_2(\Omega)$ and $L_2(\Gamma)$ respectively.

In the Sobolev case of $\varphi(t) \equiv 1$, Proposition 1 is proved in [7] (Theorem 3.4.1) for integer s and in [10] (Theorems 2.4.1 and 2.7.3) for real s and general elliptic systems. For arbitrary $\varphi \in \mathcal{M}$ and real $s > 2q$, this proposition is proved in [30] (Theorem 1). The case where $\varphi \in \mathcal{M}$ and $2q - 1/2 < s \leq 2q$ is covered in [37] (Theorem 1).

In connection with Proposition 1, we recall the definition of a Fredholm operator and its index. A linear bounded operator $T: E_1 \rightarrow E_2$ acting between certain Banach spaces E_1 and E_2 is said to be Fredholm if its kernel $\ker T$ and co-kernel $E_2/T(E_1)$ are finite-dimensional. If this operator is Fredholm, then its range $T(E_1)$ is closed in E_2 ; see, e.g., [36] (Lemma 19.1.1). The number $\text{ind } T := \dim \ker T - \dim(E_2/T(E_1))$ is called the index of the Fredholm operator T .

Proposition 1 is not true if $s \leq 2q - 1/2$. This is caused by the fact that the boundary differential operators B_j cannot be reasonably applied to all distributions $u \in H^{s,\varphi}(\Omega)$ in this case. Namely, the mapping $u \mapsto B_j u$, with $u \in C^\infty(\bar{\Omega})$, cannot be extended to a continuous operator from the whole $H^{s,\varphi}(\Omega)$ to $\mathcal{D}'(\Gamma)$ if $s \leq m_j + 1/2$.

To obtain a version of Theorem 1 for arbitrary $s \leq 2q - 1/2$, we restrict ourselves to the solutions $u \in H^{s,\varphi}(\Omega)$ of the elliptic equation $Au = f$ with $f \in L_2(\Omega) = H^0(\Omega)$. This approach originates from J.-L. Lions and E. Magenes [27, 28], who took it to the investigation of regular elliptic boundary-value problems in Sobolev spaces.

Let $s < 2q$ and $\varphi \in \mathcal{M}$. Consider the linear space

$$H_A^{s,\varphi}(\Omega) := \{u \in H^{s,\varphi}(\Omega) : Au \in L_2(\Omega)\}$$

endowed with the graph norm

$$\|u\|_{H_A^{s,\varphi}(\Omega)} := \left(\|u\|_{H^{s,\varphi}(\Omega)}^2 + \|Au\|_{L_2(\Omega)}^2 \right)^{1/2}. \tag{13}$$

Here, Au is understood in the sense of the theory of distributions.

This space is Hilbert with respect to the norm (13). Indeed, this norm is generated by an inner product because so are the norms on the right-hand side of (13). Besides, the space $H_A^{s,\varphi}(\Omega)$ is complete with respect to this norm. In fact, if (u_k) is a Cauchy sequence in this space, then there exist the limits $u := \lim u_k$ in $H^{s,\varphi}(\Omega)$ and $f := \lim Au_k$ in $L_2(\Omega)$ because the spaces $H^{s,\varphi}(\Omega)$ and $L_2(\Omega)$ are complete. The differential operator A is continuous in $\mathcal{S}'(\Omega)$, so $Au = \lim Au_k = f$ in $\mathcal{S}'(\Omega)$. Here, recall, $u \in H^{s,\varphi}(\Omega)$ and $f \in L_2(\Omega)$. Therefore, $u \in H_A^{s,\varphi}(\Omega)$ and $\lim u_k = u$ in the space $H_A^{s,\varphi}(\Omega)$. Thus, this space is complete.

It is useful to note that even when all coefficients of A are constant, the space $H_A^{s,\varphi}(\Omega)$ depends essentially on each of them. This follows from the result by L. Hörmander [38] (Theorem 3.1).

Consider the Hilbert spaces

$$\begin{aligned} \mathcal{D}_A^{s,\varphi}(\Omega, \Gamma) &:= H_A^{s,\varphi}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{s+r_k-1/2,\varphi}(\Gamma), \\ \mathcal{E}_{0,s,\varphi}(\Omega, \Gamma) &:= L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\infty} H^{s-m_j-1/2,\varphi}(\Gamma). \end{aligned}$$

We investigate the properties of the elliptic boundary-value problem (1), (2) in the couple of these spaces.

Theorem 1. *Let $s < 2q$ and $\varphi \in \mathcal{M}$. Then the set $C^\infty(\bar{\Omega})$ is dense in the space $H_A^{s,\varphi}(\Omega)$, and the mapping (3) extends uniquely (by continuity) to a bounded operator*

$$\Lambda : \mathcal{D}_A^{s,\varphi}(\Omega, \Gamma) \rightarrow \mathcal{E}_{0,s,\varphi}(\Omega, \Gamma). \tag{14}$$

This operator is Fredholm. Its kernel coincides with N , whereas its range consists of all vectors

$$(f, g) := (f, g_1, \dots, g_{q+\varkappa}) \in \mathcal{E}_{0,s,\varphi}(\Omega, \Gamma) \quad (15)$$

that satisfy condition (12). The index of the operator (14) is equal to $\dim N - \dim N^+$ and does not depend on s and φ .

If $N = \{0\}$ and $N^+ = \{0\}$, then the operator (14) is an isomorphism between the spaces $\mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)$ and $\mathcal{E}_{0,s,\varphi}(\Omega, \Gamma)$. Generally, this operator naturally induces an isomorphism

$$\Lambda : \mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)/N \leftrightarrow \mathcal{R}_{0,s,\varphi}(\Omega, \Gamma). \quad (16)$$

Here,

$$\mathcal{R}_{0,s,\varphi}(\Omega, \Gamma) := \{(f, g_1, \dots, g_{q+\varkappa}) \in \mathcal{E}_{0,s,\varphi}(\Omega, \Gamma) : (12) \text{ is true}\}$$

is a subspace of $\mathcal{E}_{0,s,\varphi}(\Omega, \Gamma)$.

Let us discuss the properties of generalized solutions to the elliptic boundary-value problem (1), (2). We first give the definition of such solutions.

Let us denote

$$\mathcal{S}'_A(\Omega) := \{u \in \mathcal{S}'(\Omega) : Au \in L_2(\Omega)\}.$$

We interpret $\mathcal{S}'_A(\Omega)$ as a subspace of the linear topological space $\mathcal{S}'(\Omega)$. Since the domain Ω is bounded, we conclude that $\mathcal{S}'_A(\Omega)$ is the union of all spaces $H_A^{s,\varphi}(\Omega)$ with $s < 2q$ and $\varphi \in \mathcal{M}$.

Consider an arbitrary vector

$$(u, v) := (u, v_1, \dots, v_\varkappa) \in \mathcal{S}'_A(\Omega) \times (\mathcal{D}'(\Gamma))^\varkappa. \quad (17)$$

Since $(u, v) \in \mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)$ for certain $s < 2q$ and $\varphi \in \mathcal{M}$, we define its image (15) by the formula $\Lambda(u, v) = (f, g)$ if we apply the bounded operator (14). The vector (u, v) does not depend on s and φ and is called a generalized solution of the boundary-value problem (1), (2) with the right-hand side (15). This solution satisfies the following a priori estimate.

Theorem 2. *Let $s < 2q$, $\varphi \in \mathcal{M}$, and $\sigma > 0$. Then there exists a number $c > 0$ such that*

$$\|(u, v)\|_{\mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)} \leq c \left(\|\Lambda(u, v)\|_{\mathcal{E}_{0,s,\varphi}(\Omega, \Gamma)} + \|(u, v)\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega, \Gamma)} \right) \quad (18)$$

for an arbitrary vector $(u, v) \in \mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)$. Here, $c = c(s, \varphi, \sigma)$ is independent of (u, v) .

If $N = \{0\}$, then the second term on the right-hand side of the inequality (18) is absent. This immediately follows from the isomorphism (16).

In connection with this theorem, we mention Peetre's lemma [39] (Lemma 3) and remark that the embedding $\mathcal{D}_A^{s,\varphi}(\Omega, \Gamma) \hookrightarrow \mathcal{D}_A^{s-\sigma,\varphi}(\Omega, \Gamma)$, with $\sigma > 0$, is not compact.

The next theorem says about the regularity properties of generalized solutions to the elliptic boundary-value problem (1), (2).

Theorem 3. *Let $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. Suppose that the vector (17) is a generalized solution to the elliptic boundary-value problem (1), (2) whose right-hand side meets the conditions*

$$f \in H^{s-2q,\varphi}(\Omega) \quad \text{if } s \geq 2q, \quad (19)$$

$$g_j \in H^{s-m_j-1/2,\varphi}(\Gamma) \quad \text{for each } j \in \{1, \dots, q + \varkappa\}. \quad (20)$$

Then the solution satisfies the conditions

$$u \in H^{s,\varphi}(\Omega), \quad (21)$$

$$v_k \in H^{s+r_k-1/2,\varphi}(\Gamma) \quad \text{for each } k \in \{1, \dots, \varkappa\}. \quad (22)$$

We conclude that the solution inherits the (supplementary) regularity φ of the right-hand sides of the elliptic problem.

We note that the restriction $s \geq 2q$ in condition (19) is caused by the following: if $s < 2q$, then

$$f = Au \in L_2(\Omega) \subset H^{s-2q,\varphi}(\Omega)$$

in view of (17) and (9). Thus, the assumption $f \in H^{s-2q,\varphi}(\Omega)$ is superfluous in Theorem 3 in the case of $s < 2q$.

5. Auxiliary results relating to the problem under consideration.. Our proof of Theorem 1 will be based on the result by V. A. Kozlov, V. G. Maz'ya and J. Rossmann [7] (Theorem 3.4.1). They demonstrated that the elliptic boundary-value problem (1), (2) is Fredholm in the two-sided scale of function spaces introduced by Ya. A. Roitberg [11, 12]. Being a certain modification of the classical Sobolev scale, this two-sided scale proves to be useful in the theory of elliptic boundary-value problems in the situation where their right-hand sides can be nonregular distributions (see monographs by Yu. M. Berezansky [40] (Chapt. III, Sect. 6) and Ya. A. Roitberg [10, 13]).

Let us recall the definition of the inner product spaces $H^{s,(2q)}(\Omega)$ introduced by Ya. A. Roitberg. For our purposes, we restrict ourselves to the case of $s \in \mathbb{Z}$.

Beforehand, we need to define the space $H^{s,(0)}(\Omega)$. If $s \geq 0$, then $H^{s,(0)}(\Omega) := H^s(\Omega)$ is the inner product Sobolev space of order s over Ω . But, if $s < 0$, then $H^{s,(0)}(\Omega)$ is the completion of $C^\infty(\bar{\Omega})$ with respect to the Hilbert norm

$$\|u\|_{H^{s,(0)}(\Omega)} := \sup \left\{ \frac{|(u, w)_\Omega|}{\|w\|_{H^{-s}(\Omega)}} : w \in H^{-s}(\Omega), w \neq 0 \right\}.$$

Thus, we have the rigging

$$H^{s,(0)}(\Omega) \hookrightarrow L_2(\Omega) \hookrightarrow H^{-s,(0)}(\Omega), \quad \text{with } s > 0,$$

of the space $L_2(\Omega)$ with positive and negative Sobolev spaces (see [40] (Chapt. I, Sect. 3, Subsect. 1)). Note that the negative space $H^{s,(0)}(\Omega)$, with $s < 0$, admits the following description: the mapping $u \mapsto \mathcal{O}u$, with $u \in C^\infty(\bar{\Omega})$, extends uniquely (by continuity) to an isometric isomorphism of the space $H^{s,(0)}(\Omega)$ onto the subspace $\{w \in H^s(\mathbb{R}^n) : \text{supp } w \subseteq \bar{\Omega}\}$ of $H^s(\mathbb{R}^n)$. Here, $\mathcal{O}u := u$ on $\bar{\Omega}$, and $\mathcal{O}u := 0$ on $\mathbb{R}^n \setminus \bar{\Omega}$.

We can now define the Hilbert space $H^{s,(2q)}(\Omega)$, with $s \in \mathbb{Z}$. By definition, the space $H^{s,(2q)}(\Omega)$ is the completion of the set $C^\infty(\bar{\Omega})$ with respect to the Hilbert norm

$$\|u\|_{H^{s,(2q)}(\Omega)} := \left(\|u\|_{H^{s,(0)}(\Omega)}^2 + \sum_{k=1}^{2q} \|(D_\nu^{k-1}u) \upharpoonright \Gamma\|_{H^{s-k+1/2}(\Gamma)}^2 \right)^{1/2}.$$

This space admits the following description [13] (Sect. 2.2): the linear mapping

$$T_{2q} : u \mapsto (u, u \upharpoonright \Gamma, \dots, (D_\nu^{2q-1} u) \upharpoonright \Gamma), \quad \text{with } u \in C^\infty(\overline{\Omega}),$$

extends uniquely (by continuity) to an isometric operator

$$T_{2q} : H^{s,(2q)}(\Omega) \rightarrow H^{s,(0)}(\Omega) \oplus \bigoplus_{k=1}^{2q} H^{s-k+1/2}(\Gamma) =: \Pi_{s,(2q)}(\Omega, \Gamma);$$

the range of this operator consists of all vectors

$$(u_0, u_1, \dots, u_{2q}) \in \Pi_{s,(2q)}(\Omega, \Gamma)$$

such that $u_k = R_\Gamma D_\nu^{k-1} u_0$ for all integers $k \in \{1, \dots, 2q\}$ satisfying $s > k - 1/2$. Thus, we can interpret an arbitrary element $u \in H^{s,(2q)}(\Omega)$ as the vector $(u_0, u_1, \dots, u_{2q}) := T_{2q} u$. This interpretation is taken in the mentioned monograph [7] (Sect. 3.2.1) as a definition of the space $H^{s,(2q)}(\Omega)$.

Note that

$$H^{s,(2q)}(\Omega) = H^s(\Omega) \quad \text{for all } s \geq 2q \quad (23)$$

as completions of $C^\infty(\overline{\Omega})$ with respect to equivalent norms.

We can now formulate the result by V. A. Kozlov, V. G. Maz'ya and J. Rossmann [7] (Theorem 3.4.1). Consider the Hilbert spaces

$$\mathcal{D}^{s,(2q)}(\Omega, \Gamma) := H^{s,(2q)}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{s+r_k-1/2}(\Gamma),$$

$$\mathcal{E}^{s-2q,(0)}(\Omega, \Gamma) := H^{s-2q,(0)}(\Omega) \oplus \bigoplus_{j=1}^{q+\infty} H^{s-m_j-1/2}(\Gamma),$$

with $s \in \mathbb{Z}$.

Proposition 2. *For arbitrary $s \in \mathbb{Z}$, the mapping (3) extends uniquely (by continuity) to a bounded operator*

$$\Lambda : \mathcal{D}^{s,(2q)}(\Omega, \Gamma) \rightarrow \mathcal{E}^{s-2q,(0)}(\Omega, \Gamma). \quad (24)$$

This operator is Fredholm. Its kernel coincides with N , and its range consists of all vectors

$$(f, g_1, \dots, g_{q+\infty}) \in \mathcal{E}^{s-2q,(0)}(\Omega, \Gamma) \quad (25)$$

that satisfy (12). The index of the operator (24) is equal to $\dim N - \dim N^+$ and does not depend on s .

Note that if $s \in \mathbb{Z}$, $s \geq 2q$, and $\varphi \equiv 1$, then Proposition 1 coincides with Proposition 2 in view of (23).

In the proof of Theorem 1, we will use Proposition 2 together with the following result. For an arbitrary integer $s < 2q$, we consider the linear space

$$H_A^{s,(2q)}(\Omega) := \{u \in H^{s,(2q)}(\Omega) : Au \in L_2(\Omega)\}$$

endowed with the graph norm

$$\|u\|_{H_A^{s,(2q)}(\Omega)} := \left(\|u\|_{H^{s,(2q)}(\Omega)}^2 + \|Au\|_{L_2(\Omega)}^2 \right)^{1/2}. \quad (26)$$

Note that the image $Au \in H^{s-2q,(0)}(\Omega)$ is well defined for each $u \in H^{s,(2q)}(\Omega)$ by closure because the mapping $u \mapsto Au$, with $u \in C^\infty(\overline{\Omega})$, extends by continuity to a bounded operator $A: H^{s,(2q)}(\Omega) \rightarrow H^{s-2q,(0)}(\Omega)$ for every $s \in \mathbb{Z}$ [13] (Lemma 2.3.1). The space $H_A^{s,(2q)}(\Omega)$ is Hilbert with respect to the norm (26), and the set $C^\infty(\overline{\Omega})$ is dense in it (see, e.g., [23] (Sect. 4.4.2, the proof of Theorem 4.25)).

Proposition 3. *For an arbitrary integer $s < 2q$, the identity mapping on $C^\infty(\overline{\Omega})$ extends uniquely (by continuity) to an isomorphism between the spaces $H_A^{s,(2q)}(\Omega)$ and $H_A^s(\Omega)$. Thus, these spaces are equal as completions of the set $C^\infty(\overline{\Omega})$ with respect to equivalent norms.*

Here and below, $H_A^s(\Omega)$ is the space $H_A^{s,\varphi}(\Omega)$ in the Sobolev case of $\varphi \equiv 1$.

The proof of Proposition 3 is given in monograph [23] (Sect. 4.4.2, see the reasoning from formula (4.196) to the end of the proof of Theorem 4.25).

6. Auxiliary results relating to interpolation between Hilbert spaces.. The refined Sobolev scales possess an important interpolation property with respect to inner product Sobolev spaces. Namely, each space $H^{s,\varphi}(G)$, with $s \in \mathbb{R}$, $\varphi \in \mathcal{M}$, and $G \in \{\mathbb{R}^n, \Omega, \Gamma\}$, can be obtained by the interpolation with an appropriate function parameter between the Sobolev spaces $H^{s-\varepsilon}(G)$ and $H^{s+\delta}(G)$ with $\varepsilon, \delta > 0$. This property will play a key role in our proof of Theorem 1 for arbitrary $\varphi \in \mathcal{M}$. Therefore, we recall the definition of this interpolation in the case of arbitrary Hilbert spaces and discuss some of its properties that are necessary for us.

The method of interpolation with a function parameter between Hilbert spaces appeared first in C. Foias and J.-L. Lions' paper [41, p. 278]. We follow monograph [23] (Sect. 1.1), which systematically expounds this method. For our purposes, it is sufficient to restrict ourselves to separable Hilbert spaces.

Let $X := [X_0, X_1]$ be a given ordered couple of separable complex Hilbert spaces X_0 and X_1 such that the continuous and dense embedding $X_1 \hookrightarrow X_0$ holds. This couple is said to be admissible. For X , there exists a self-adjoint positive-definite operator J on X_0 that has the domain X_1 and that satisfies the condition $\|Jw\|_{X_0} = \|w\|_{X_1}$ for each $w \in X_1$. This operator is uniquely determined by the couple X and is called a generating operator for X . It defines an isometric isomorphism $J: X_1 \leftrightarrow X_0$.

Let \mathcal{B} denote the set of all Borel measurable functions $\psi: (0, \infty) \rightarrow (0, \infty)$ such that ψ is bounded on each compact interval $[a, b]$, with $0 < a < b < \infty$, and that $1/\psi$ is bounded on every set $[r, \infty)$, with $r > 0$.

For arbitrary $\psi \in \mathcal{B}$, we consider the (generally, unbounded) operator $\psi(J)$, which is defined on X_0 as the Borel function ψ of J and is built with the help of Spectral Theorem applied to the self-adjoint operator J . Let $[X_0, X_1]_\psi$ or, simply, X_ψ denote the domain of the operator $\psi(J)$ endowed with the norm $\|u\|_{X_\psi} = \|\psi(J)u\|_{X_0}$. The space X_ψ is Hilbert and separable with respect to this norm.

A function $\psi \in \mathcal{B}$ is called an interpolation parameter if the following condition is fulfilled for all admissible couples $X = [X_0, X_1]$ and $Y = [Y_0, Y_1]$ of Hilbert spaces and for an arbitrary linear mapping T given on X_0 : if the restriction of T to X_j is a bounded operator $T: X_j \rightarrow Y_j$ for each $j \in \{0, 1\}$, then the restriction of T to X_ψ is also a bounded operator $T: X_\psi \rightarrow Y_\psi$. In this case we say that X_ψ is obtained by the interpolation with the function parameter ψ of the couple X (or between the spaces X_0 and X_1).

The function $\psi \in \mathcal{B}$ is an interpolation parameter if and only if ψ is pseudoconcave on a neighbourhood of infinity, i.e., $\psi(t) \asymp \psi_1(t)$ with $t \gg 1$ for a certain positive concave function $\psi_1(t)$. (As usual, $\psi \asymp \psi_1$ means that both the functions ψ/ψ_1 and ψ_1/ψ are bounded on the indicated set). This fundamental property of the interpolation follows from J. Peetre's results [42, 43]. Specifically, if $\psi(t) \equiv t^\sigma \psi_0(t)$ for some number $\sigma \in (0, 1)$ and function ψ_0 varying slowly at infinity (in the sense of J. Karamata), then ψ is an interpolation parameter.

Let us formulate the above-mentioned interpolation property of the refined Sobolev scales [23] (Theorems 1.14, 2.2, and 3.2).

Proposition 4. *Let a function $\varphi \in \mathcal{M}$ and positive real numbers ε, δ be given. Define a function $\psi \in \mathcal{B}$ by the formula*

$$\psi(t) = \begin{cases} t^{\varepsilon/(\varepsilon+\delta)} \varphi(t^{1/(\varepsilon+\delta)}) & \text{if } t \geq 1, \\ \varphi(1) & \text{if } 0 < t < 1. \end{cases} \quad (27)$$

Then ψ is an interpolation parameter, and

$$[H^{s-\varepsilon}(G), H^{s+\delta}(G)]_\psi = H^{s,\varphi}(G) \quad \text{for each } s \in \mathbb{R}$$

with equivalence of norms. Here, $G \in \{\mathbb{R}^n, \Omega, \Gamma\}$. If $G = \mathbb{R}^n$, then the equality of norms holds.

We will also use the following three general properties of the interpolation.

Proposition 5. *Let $X = [X_0, X_1]$ and $Y = [Y_0, Y_1]$ be admissible couples of Hilbert spaces, and let a linear mapping T be given on X_0 . Suppose that we have the bounded and Fredholm operators $T: X_j \rightarrow Y_j$, with $j = 0, 1$, that possess the common kernel and the common index. Then, for an arbitrary interpolation parameter $\psi \in \mathcal{B}$, the bounded operator $T: X_\psi \rightarrow Y_\psi$ is Fredholm, has the same kernel and the same index, and, moreover, its range $T(X_\psi) = Y_\psi \cap T(X_0)$.*

The proof of this proposition is given in [23] (Sect. 1.1.7).

Proposition 6. *Let $[X_0^{(j)}, X_1^{(j)}]$, with $j = 1, \dots, r$, be a finite collection of admissible couples of Hilbert spaces. Then, for every function $\psi \in \mathcal{B}$, we have*

$$\left[\bigoplus_{j=1}^r X_0^{(j)}, \bigoplus_{j=1}^r X_1^{(j)} \right]_\psi = \bigoplus_{j=1}^r [X_0^{(j)}, X_1^{(j)}]_\psi$$

with equality of norms.

The proof of this proposition is given in [23] (Sect. 1.1.5).

The third property deals with the interpolation of subspaces connected with a linear bounded operator. Beforehand, we admit the following notation.

Let H, Φ and Ψ be Hilbert spaces, with $\Phi \hookrightarrow \Psi$ continuously, and let a bounded linear operator $T: H \rightarrow \Psi$ be given. We put

$$(H)_{T,\Phi} := \{u \in H : Tu \in \Phi\}.$$

The linear space $(H)_{T,\Phi}$ is considered as a Hilbert space with respect to the graph norm

$$\|u\|_{(H)_{T,\Phi}} := (\|u\|_H^2 + \|Tu\|_\Phi^2)^{1/2}.$$

Note that this space does not depend on Ψ .

Proposition 7. *Suppose that six separable Hilbert spaces X_0, Y_0, Z_0, X_1, Y_1 , and Z_1 and three linear mappings T, R , and S satisfy the following seven conditions:*

- (i) the couples $X = [X_0, X_1]$ and $Y = [Y_0, Y_1]$ are admissible;
 - (ii) the spaces Z_0 and Z_1 are subspaces of a certain linear space E ;
 - (iii) the continuous embeddings $Y_0 \hookrightarrow Z_0$ and $Y_1 \hookrightarrow Z_1$ hold;
 - (iv) the mapping T is given on X_0 and defines the bounded operators $T : X_0 \rightarrow Z_0$ and $T : X_1 \rightarrow Z_1$;
 - (v) the mapping R is given on E and defines the bounded operators $R : Z_0 \rightarrow X_0$ and $R : Z_1 \rightarrow X_1$;
 - (vi) the mapping S is given on E and defines the bounded operators $S : Z_0 \rightarrow Y_0$ and $S : Z_1 \rightarrow Y_1$;
 - (vii) the equality $TR\omega = \omega + S\omega$ holds for each $\omega \in E$.
- Then the couple of spaces $[(X_0)_{T,Y_0}, (X_1)_{T,Y_1}]$ is admissible, and

$$[(X_0)_{T,Y_0}, (X_1)_{T,Y_1}]_\psi = (X_\psi)_{T,Y_\psi}$$

up to equivalence of norms for an arbitrary interpolation parameter $\psi \in \mathcal{B}$.

An analog of Proposition 7 appeared first in J.-L. Lions and E. Magenes' monograph [44] (Theorem 14.3), where the case of holomorphic interpolation was considered. This proposition is proved by V. A. Mikhailets and A. A. Murach in paper [18] (Sect. 4); see also monograph [23] (Theorem 3.12).

7. Proofs of the main results. Let us prove Theorems 1–3.

Proof of Theorem 1. We will first prove this theorem in the Sobolev case where $s \in \mathbb{Z}$, $s < 2q$, and $\varphi \equiv 1$. Consider the bounded and Fredholm operator (24) from Proposition 2. Certainly, the restriction of this operator to the space $H_A^{s,(2q)}(\Omega)$ is a bounded operator

$$\Lambda : H_A^{s,(2q)}(\Omega) \oplus \bigoplus_{k=1}^{\varkappa} H^{s+r_k-1/2}(\Gamma) \rightarrow L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{s-m_j-1/2}(\Gamma). \tag{28}$$

Since $C^\infty(\overline{\Omega})$ is dense in $H_A^{s,(2q)}(\Omega)$, this operator is an extension by continuity of the mapping (3). Let $\Lambda_{s,(2q)}$ denote the operator (24), and let Λ_s denote the operator (28). Evidently, $\ker \Lambda_s = \ker \Lambda_{s,(2q)}$ and

$$\text{Ran } \Lambda_s = \{(f, g_1, \dots, g_{q+\varkappa}) \in \text{Ran } \Lambda_{s,(2q)} : f \in L_2(\Omega)\},$$

with Ran denoting the range of the corresponding operator. It follows from this by Proposition 2 that $\ker \Lambda_s = N$ and

$$\text{Ran } \Lambda_s = \left\{ (f, g_1, \dots, g_{q+\varkappa}) \in L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{s-m_j-1/2}(\Gamma) : (12) \text{ is true} \right\}.$$

Therefore,

$$\dim \ker \Lambda_s = \dim N < \infty \quad \text{and} \quad \dim \text{coker } \Lambda_s = \dim N^+ < \infty.$$

Thus, the bounded operator Λ_s is Fredholm. In view of Proposition 3, we conclude that Λ_s is the required operator (14) in Theorem 1, the set $C^\infty(\overline{\Omega})$ being dense in $H_A^s(\Omega)$.

We will now prove Theorem 1 in the general situation with the help of interpolation with a function parameter between Sobolev spaces. Let $s \in \mathbb{R}$, $s < 2q$, and $\varphi \in \mathcal{M}$. Choose an integer $p \geq 2$ such that $s > -2q(p-1)$. The mapping (3) extends by continuity to the bounded and Fredholm operators

$$\Lambda: H_A^{-2q(p-1)}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{-2q(p-1)+r_k-1/2}(\Gamma) \rightarrow L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\infty} H^{-2q(p-1)-m_j-1/2}(\Gamma) \quad (29)$$

and

$$\Lambda: H^{2q}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{2q+r_k-1/2}(\Gamma) \rightarrow L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\infty} H^{2q-m_j-1/2}(\Gamma). \quad (30)$$

This has just been proved for the first operator and is said in Proposition 2 for the second. These operators have the common kernel N and the same index equal to $\dim N - \dim N^+$. The range of the first operator satisfies the equation

$$\begin{aligned} & \Lambda(H_A^{-2q(p-1)}(\Omega)) = \\ & = \left\{ (f, g_1, \dots, g_{q+\infty}) \in L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\infty} H^{-2q(p-1)-m_j-1/2}(\Gamma) : (12) \text{ is true} \right\}, \quad (31) \end{aligned}$$

and an analogous formula holds for the second operator. Note that the second operator is a restriction of the first.

Let us define the interpolation parameter ψ by formula (27) in which $\varepsilon := s + 2q(p-1) > 0$ and $\delta := 2q - s > 0$. Applying the interpolation with the function parameter ψ to the couples of spaces in (29) and (30), we obtain the bounded operator

$$\begin{aligned} \Lambda: & \left[H_A^{-2q(p-1)}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{-2q(p-1)+r_k-1/2}(\Gamma), H^{2q}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{2q+r_k-1/2}(\Gamma) \right]_{\psi} \rightarrow \\ & \rightarrow \left[L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\infty} H^{-2q(p-1)-m_j-1/2}(\Gamma), L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\infty} H^{2q-m_j-1/2}(\Gamma) \right]_{\psi}. \quad (32) \end{aligned}$$

According to Proposition 5, this operator is Fredholm with kernel N and index $\dim N - \dim N^+$.

Let us describe the interpolation spaces in (32). Using Propositions 6 and 4 successively, we get

$$\begin{aligned} & \left[H_A^{-2q(p-1)}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{-2q(p-1)+r_k-1/2}(\Gamma), H^{2q}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{2q+r_k-1/2}(\Gamma) \right]_{\psi} = \\ & = [H_A^{-2q(p-1)}(\Omega), H^{2q}(\Omega)]_{\psi} \oplus \bigoplus_{k=1}^{\infty} [H^{-2q(p-1)+r_k-1/2}(\Gamma), H^{2q+r_k-1/2}(\Gamma)]_{\psi} = \\ & = [H_A^{-2q(p-1)}(\Omega), H^{2q}(\Omega)]_{\psi} \oplus \bigoplus_{k=1}^{\infty} H^{s+r_k-1/2, \varphi}(\Gamma). \quad (33) \end{aligned}$$

Here, we take into account the equalities

$$s - \varepsilon = -2q(p - 1) \quad \text{and} \quad s + \delta = 2q. \tag{34}$$

Analogously,

$$\left[L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{-2q(p-1)-m_j-1/2}(\Gamma), L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{2q-m_j-1/2}(\Gamma) \right]_{\psi} = L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{s-m_j-1/2,\varphi}(\Gamma). \tag{35}$$

These equalities of spaces hold true up to equivalence of norms.

Let us prove that

$$[H_A^{-2q(p-1)}(\Omega), H^{2q}(\Omega)]_{\psi} = H_A^{s,\varphi}(\Omega) \tag{36}$$

up to equivalence of norms. To this end, we use Proposition 7 in which

$$X_0 := H^{-2q(p-1)}(\Omega), \quad X_1 := H^{2q}(\Omega), \quad Y_0 := Y_1 := Z_1 := L_2(\Omega), \quad Z_0 := E := H^{-2qp}(\Omega),$$

and $T := A$. Then

$$H_A^{-2q(p-1)}(\Omega) = (X_0)_{T,Y_0} \quad \text{and} \quad H^{2q}(\Omega) = (X_1)_{T,Y_1}. \tag{37}$$

Note that the latter equality holds true up to equivalence of norms because A is a bounded operator from $H^\sigma(\Omega)$ to $H^{\sigma-2q}(\Omega)$ for each $\sigma \in \mathbb{R}$, specifically, from $H^{2q}(\Omega)$ to $L_2(\Omega)$.

Evidently, conditions (i)–(iv) of Proposition 7 are fulfilled. We also need to give certain operators R and S satisfying conditions (v)–(vii). With this in mind, we use the known fact that the mapping $u \mapsto A^p A^{p+} u + u$ defines the isomorphism

$$A^p A^{p+} + I : H_D^\sigma(\Omega) \leftrightarrow H^{\sigma-4qp}(\Omega) \quad \text{for each } \sigma \geq 2qp \tag{38}$$

(see, e.g., [23] (Lemma 3.1)). Here, of course, A^p is the p -th iteration of A , then A^{p+} is the formally adjoint operator to the differential operator A^p , and I is the identity operator. Besides,

$$H_D^\sigma(\Omega) := \{u \in H^\sigma(\Omega) : R_\Gamma D_\nu^{j-1} u = 0 \text{ for each } j \in \{1, \dots, 2qp\}\}$$

is a subspace of $H^\sigma(\Omega)$. The inverse of (38) defines the bounded linear operator

$$(A^p A^{p+} + I)^{-1} : H^\sigma(\Omega) \rightarrow H^{\sigma+4qp}(\Omega) \quad \text{for each } \sigma \geq -2qp. \tag{39}$$

We now let

$$R := A^{p-1} A^{p+} (A^p A^{p+} + I)^{-1} \quad \text{and} \quad S = -(A^p A^{p+} + I)^{-1}.$$

Using (39), we get the bounded operators

$$R : Z_0 = H^{-2qp}(\Omega) \rightarrow H^{2qp-2qp-2q(p-1)}(\Omega) = X_0,$$

$$R: Z_1 = L_2(\Omega) \rightarrow H^{4qp-2qp-2q(p-1)}(\Omega) = X_1,$$

$$S: Z_0 = H^{-2qp}(\Omega) \rightarrow H^{2qp}(\Omega) \hookrightarrow L_2(\Omega) = Y_0,$$

$$S: Z_1 = L_2(\Omega) \rightarrow H^{4qp}(\Omega) \hookrightarrow L_2(\Omega) = Y_1;$$

here, the embeddings are continuous. Besides,

$$AR = AA^{p-1}A^{p+}(A^pA^{p+} + I)^{-1} = (A^pA^{p+} + I - I)(A^pA^{p+} + I)^{-1} = I + S$$

on $E = H^{-2qp}(\Omega)$. Thus, the rest conditions (v)–(vii) of Proposition 7 are satisfied.

Now, according to this proposition and in view of (37), we get

$$[H_A^{-2q(p-1)}(\Omega), H^{2q}(\Omega)]_\psi = [(X_0)_{T,Y_0}, (X_1)_{T,Y_1}]_\psi = (X_\psi)_{T,Y_\psi}.$$

Here, by Proposition 4 and (34), we have

$$X_\psi = [H^{-2q(p-1)}(\Omega), H^{2q}(\Omega)]_\psi = H^{s,\varphi}(\Omega).$$

Besides,

$$Y_\psi = [L_2(\Omega), L_2(\Omega)]_\psi = L_2(\Omega).$$

These three formulas immediately gives the required equality (36).

Note that

$$\text{the set } C^\infty(\bar{\Omega}) \text{ is dense in the space } H_A^{s,\varphi}(\Omega). \quad (40)$$

This follows from the density of $C^\infty(\bar{\Omega})$ in $H^{2q}(\Omega)$ and from the dense continuous embedding of $H^{2q}(\Omega)$ in $H_A^{s,\varphi}(\Omega)$. The latter embedding is a general property of the interpolation used in (36).

It follows from the interpolation formulas (33), (35), and (36) that the bounded and Fredholm operator (32) acts between the spaces

$$\Lambda: H_A^{s,\varphi}(\Omega) \oplus \bigoplus_{k=1}^{\varkappa} H^{s+r_k-1/2,\varphi}(\Gamma) \rightarrow L_2(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{s-m_j-1/2,\varphi}(\Gamma). \quad (41)$$

(Recall that these spaces are denoted by $\mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)$ and $\mathcal{E}_{0,s,\varphi}(\Omega, \Gamma)$ respectively.) We have proved that the operator (41) has the kernel N and the index $\dim N - \dim N^+$. Furthermore, according to Proposition 5 and formula (31), the range of this operator is equal to

$$\mathcal{E}_{0,s,\varphi}(\Omega, \Gamma) \cap \Lambda(H_A^{-2q(p-1)}(\Omega)) = \{(f, g_1, \dots, g_{q+\varkappa}) \in \mathcal{E}_{0,s,\varphi}(\Omega, \Gamma) : (12) \text{ is true}\}.$$

Finally, in view of (40), the operator (41) is an extension by continuity of the mapping (3). Thus, this operator is the required operator (14).

Theorem 1 is proved.

Proof of Theorem 2. It follows from the isomorphism (16) that

$$\inf \{ \|(u, v) + (u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)} : (u^{(0)}, v^{(0)}) \in N \} \leq c_0 \|\Lambda(u, v)\|_{\mathcal{E}_{0,s,\varphi}(\Omega, \Gamma)} \quad (42)$$

for each $(u, v) \in \mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)$, with c_0 being the norm of the inverse operator to (16). Since N is a finite-dimensional subspace of both the spaces $\mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)$ and $\mathcal{D}_A^{s-\sigma,\varphi}(\Omega, \Gamma)$, the norms in them are

equivalent on N . Hence, for each $(u^{(0)}, v^{(0)}) \in N$, we can write

$$\|(u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma)} \leq c_1 \|(u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)}$$

with some number $c_1 > 0$, which is independent of (u, v) and $(u^{(0)}, v^{(0)})$. Besides,

$$\begin{aligned} \|(u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)} &\leq \|(u, v) + (u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)} + \|(u, v)\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)} \leq \\ &\leq c_2 \|(u, v) + (u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma)} + \|(u, v)\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)}. \end{aligned}$$

Here, c_2 is the norm of the continuous embedding operator $\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma) \hookrightarrow \mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)$. Hence,

$$\begin{aligned} \|(u, v)\|_{\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma)} &\leq \|(u, v) + (u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma)} + \|(u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma)} \leq \\ &\leq \|(u, v) + (u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma)} + c_1 \|(u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)} \leq \\ &\leq (1 + c_1 c_2) \|(u, v) + (u^{(0)}, v^{(0)})\|_{\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma)} + c_1 \|(u, v)\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)}. \end{aligned}$$

Passing in this inequality to the infimum over all $(u^{(0)}, v^{(0)}) \in N$ and using the bound (42), we arrive at the estimate (18); namely, we get

$$\|(u, v)\|_{\mathcal{D}_A^{s,\varphi}(\Omega,\Gamma)} \leq (1 + c_1 c_2) c_0 \|\Lambda(u, v)\|_{\mathcal{E}_{0,s,\varphi}(\Omega,\Gamma)} + c_1 \|(u, v)\|_{\mathcal{D}_A^{s-\sigma,\varphi}(\Omega,\Gamma)}.$$

Theorem 2 is proved.

Proof of Theorem 3. We first consider the case of $s < 2q$. By virtue of Theorem 1, the vector $(f, g) = \Lambda(u, v)$ satisfies (12). Note that $(f, g) \in \mathcal{E}_{0,s,\varphi}(\Omega, \Gamma)$ by the condition (20). Therefore, according to Theorem 1, we get the inclusion $(f, g) \in \Lambda(\mathcal{D}_A^{s,\varphi}(\Omega, \Gamma))$. Thus, together with $\Lambda(u, v) = (f, g)$, we have the equality $\Lambda(u', v') = (f, g)$ for a certain vector $(u', v') \in \mathcal{D}_A^{s,\varphi}(\Omega, \Gamma)$. Hence, $\Lambda(u - u', v - v') = 0$, which implies that

$$(u - u', v - v') \in N \subset C^\infty(\bar{\Omega}) \times (C^\infty(\Gamma))^\varkappa.$$

Thus,

$$(u, v) = (u', v') + (u - u', v - v') \in \mathcal{D}_A^{s,\varphi}(\Omega, \Gamma).$$

Theorem 3 has been proved in the case of $s < 2q$.

Let us examine the case of $s \geq 2q$. According to what has just been proved, the condition

$$g_j \in H^{s-m_j-1/2,\varphi}(\Gamma) \subset H^{2q-1/4-m_j-1/2,\varphi}(\Gamma) \quad \text{for each } j \in \{1, \dots, q + \varkappa\}$$

implies the inclusion

$$(u, v) \in H^{2q-1/4,\varphi}(\Omega) \oplus \bigoplus_{k=1}^{\varkappa} H^{2q-1/4+r_k-1/2,\varphi}(\Gamma).$$

Repeating the above reasoning and using Proposition 1 instead of Theorem 1, we deduce from this inclusion and the condition of Theorem 3, that the vector (u, v) satisfies (21) and (22).

Theorem 3 is proved.

1. *Lawruk B.* Parametric boundary-value problems for elliptic systems of linear differential equations. I. Construction of conjugate problems // Bull. Acad. pol. sci. Sér. sci. math., astron. et phys. – 1963. – **11**, № 5. – P. 257–267 (in Russian).
2. *Lawruk B.* Parametric boundary-value problems for elliptic systems of linear differential equations. II. A boundary-value problem for a half-space // Bull. Acad. pol. sci. Sér. sci. math., astron. et phys. – 1963. – **11**, № 5. – P. 269–278 (in Russian).
3. *Lawruk B.* Parametric boundary-value problems for elliptic systems of linear differential equations. III. Conjugate boundary problem for a half-space // Bull. Acad. pol. sci. Sér. sci. math., astron. et phys. – 1965. – **13**, № 2. – P. 105–110 (in Russian).
4. *Aslanyan A. G., Vassiliev D. G., Lidskii V. B.* Frequences of free oscillations of thin shell interacting with fluid // Funct. Anal. and Appl. – 1981. – **15**, № 3. – P. 157–164.
5. *Ciarlet P. G.* Plates and junctions in elastic multistructures. An asymptotic analysis. – Paris: Mayson, 1990. – viii + 218 p.
6. *Nazarov S., Pileckas K.* On noncompact free boundary problems for the plane stationary Navier–Stokes equations // J. reine und angew. Math. – 1993. – **438**. – S. 103–141.
7. *Kozlov V. A., Maz'ya V. G., Rossmann J.* Elliptic boundary value problems in domains with point singularities. – Providence: Amer. Math. Soc., 1997. – 414 p.
8. *Roitberg I. Ya.* Elliptic boundary value problems for general systems of equations in complete scales of Banach spaces // Dokl. Akad. Nauk. – 1997. – **354**, № 1. – P. 25–29 (in Russian).
9. *Roitberg I. Ya.* Elliptic boundary value problems for general elliptic systems in complete scales of Banach spaces // Oper. Theory: Adv. and Appl. – 1998. – **102**. – P. 231–241.
10. *Roitberg Ya. A.* Boundary value problems in the spaces of distributions. – Dordrecht: Kluwer Acad. Publ., 1999. – x + 276 p.
11. *Roitberg Ya. A.* Elliptic problems with nonhomogeneous boundary conditions and local increase of smoothness up to the boundary for generalized solutions // Dokl. Math. – 1964. – **5**. – P. 1034–1038.
12. *Roitberg Ya. A.* A theorem on the homeomorphisms induced in L_p by elliptic operators and the local smoothing of generalized solutions // Ukr. Mat. Zh. – 1965. – **17**, № 5. – P. 122–129 (in Russian).
13. *Roitberg Ya. A.* Elliptic boundary value problems in the spaces of distributions. – Dordrecht: Kluwer Acad. Publ., 1996. – 427 p.
14. *Hörmander L.* Linear partial differential operators. – Berlin: Springer, 1963. – 285 p.
15. *Mikhailets V. A., Murach A. A.* Elliptic operators in a refined scale of function spaces // Ukr. Math. J. – 2005. – **57**, № 5. – P. 817–825.
16. *Mikhailets V. A., Murach A. A.* Refined scales of spaces and elliptic boundary-value problems. II // Ukr. Math. J. – 2006. – **58**, № 3. – P. 398–417.
17. *Mikhailets V. A., Murach A. A.* Interpolation with a function parameter and refined scale of spaces // Meth. Funct. Anal. and Top. – 2008. – **14**, № 1. – P. 81–100.
18. *Mikhailets V. A., Murach A. A.* A regular elliptic boundary value problem for a homogeneous equation in a two-sided refined scale of spaces // Ukr. Math. J. – 2006. – **58**, № 11. – P. 1748–1767.
19. *Mikhailets V. A., Murach A. A.* An elliptic operator with homogeneous regular boundary conditions in a two-sided refined scale of spaces // Ukr. Math. Bull. – 2006. – **3**, № 4. – P. 529–560.
20. *Mikhailets V. A., Murach A. A.* Refined scale of spaces and elliptic boundary-value problems. III // Ukr. Math. J. – 2007. – **59**, № 5. – P. 744–765.
21. *Mikhailets V. A., Murach A. A.* An elliptic boundary-value problem in a two-sided refined scale of spaces // Ukr. Math. J. – 2008. – **60**, № 4. – P. 574–597.
22. *Mikhailets V. A., Murach A. A.* The refined Sobolev scale, interpolation, and elliptic problems // Banach J. Math. Anal. – 2012. – **6**, № 2. – P. 211–281.
23. *Mikhailets V. A., Murach A. A.* Hörmander spaces, interpolation, and elliptic problems. – Berlin; Boston: De Gruyter, 2014. – xii + 297 p. (Russian version is available as arXiv:1106.3214.)
24. *Slenzak G.* Elliptic problems in a refined scale of spaces // Moscow Univ. Math. Bull. – 1974. – **29**, № 3–4. – P. 80–88.
25. *Anop A. V., Murach A. A.* Parameter-elliptic problems and interpolation with a function parameter // Meth. Funct. Anal. and Top. – 2014. – **20**, № 2. – P. 103–116.

26. Anop A. V., Murach A. A. Regular elliptic boundary-value problems in the extended Sobolev scale // Ukr. Math. J. – 2014. – **66**, № 7. – P. 969–985.
27. Lions J.-L., Magenes E. Problèmes aux limites non homogènes, V // Ann. Scuola norm. super. Pisa (3). – 1962. – **16**. – P. 1–44 (in Italian).
28. Lions J.-L., Magenes E. Problèmes aux limites non homogènes, VI // J. Anal. Math. – 1963. – **11**. – P. 165–188.
29. Murach A. A. Extension of some Lions–Magenes theorems // Meth. Funct. Anal. and Top. – 2009. – **15**, № 2. – P. 152–167.
30. Chepurukhina I. S. On some classes of elliptic boundary-value problems in spaces of generalized smoothness // Different. Equat. and Relat. Matters. Zb. prac' Inst. Mat. NAN Ukrayiny. – 2014. – **11**, № 2. – P. 284–304 (in Ukrainian).
31. Karamata J. Sur certains “Tauberian theorems” de M. M. Hardy et Littlewood // Mathematica (Cluj). – 1930. – **3**. – P. 33–48.
32. Seneta E. Regularly varying functions. – Berlin: Springer, 1976. – 112 p.
33. Bingham N. H., Goldie C. M., Teugels J. L. Regular variation. – Cambridge: Cambridge Univ. Press, 1989. – 512 p.
34. Hörmander L. The analysis of linear partial differential operators. II: Differential operators with constant coefficients. – Berlin: Springer, 1983. – viii + 391 p.
35. Volevich L. R., Paneah B. P. Certain spaces of generalized functions and embedding theorems // Uspekhi Mat. Nauk. – 1965. – **20**, № 1 – P. 3–74 (in Russian).
36. Hörmander L. The analysis of linear partial differential operators. III: Pseudodifferential operators. – Berlin: Springer, 1985. – viii + 525 p.
37. Chepurukhina I. S., Murach A. A. Elliptic problems in the sense of B. Lawruk on two-sided refined scale of spaces // Meth. Funct. Anal. and Top. – 2015. – **21**, № 1. – P. 6–21.
38. Hörmander L. On the theory of general partial differential equations // Acta Math. – 1955. – **94**, № 1. – P. 161–248.
39. Peetre J. Another approach to elliptic boundary problems // Commun Pure and Appl. Math. – 1961. – **14**, № 4. – P. 711–731.
40. Berezansky Yu. M. Expansions in eigenfunctions of selfadjoint operators. – Providence: Amer. Math. Soc., 1968.
41. Foias C., Lions J.-L. Sur certains théorèmes d'interpolation // Acta Sci. Math. (Szeged). – 1961. – **22**, № 3-4. – P. 269–282.
42. Peetre J. On interpolation functions // Acta Sci. Math. (Szeged). – 1966. – **27**. – P. 167–171.
43. Peetre J. On interpolation functions. II // Acta Sci. Math. (Szeged). – 1968. – **29**, № 1-2. – P. 91–92.
44. Lions J.-L., Magenes E. Non-homogeneous boundary value problems and applications. – New York: Springer, 1972. – Vol. 1. – xvi + 357 p.

Received 13.01.15